

Measurement of CP Asymmetries in $B^0 \rightarrow \phi K^0$ and $B^0 \rightarrow K^+ K^- K_S^0$ Decays

B. Aubert,¹ R. Barate,¹ D. Boutigny,¹ F. Couderc,¹ Y. Karyotakis,¹ J. P. Lees,¹ V. Poireau,¹ V. Tisserand,¹ A. Zghiche,¹ E. Grauges-Pous,² A. Palano,³ A. Pompili,³ J. C. Chen,⁴ N. D. Qi,⁴ G. Rong,⁴ P. Wang,⁴ Y. S. Zhu,⁴ G. Eigen,⁵ I. Ofte,⁵ B. Stugu,⁵ G. S. Abrams,⁶ A. W. Borgland,⁶ A. B. Breon,⁶ D. N. Brown,⁶ J. Button-Shafer,⁶ R. N. Cahn,⁶ E. Charles,⁶ C. T. Day,⁶ M. S. Gill,⁶ A. V. Gritsan,⁶ Y. Groysman,⁶ R. G. Jacobsen,⁶ R. W. Kadel,⁶ J. Kadyk,⁶ L. T. Kerth,⁶ Yu. G. Kolomensky,⁶ G. Kukartsev,⁶ G. Lynch,⁶ L. M. Mir,⁶ P. J. Oddone,⁶ T. J. Orimoto,⁶ M. Pripstein,⁶ N. A. Roe,⁶ M. T. Ronan,⁶ W. A. Wenzel,⁶ M. Barrett,⁷ K. E. Ford,⁷ T. J. Harrison,⁷ A. J. Hart,⁷ C. M. Hawkes,⁷ S. E. Morgan,⁷ A. T. Watson,⁷ M. Fritsch,⁸ K. Goetzen,⁸ T. Held,⁸ H. Koch,⁸ B. Lewandowski,⁸ M. Pelizaeus,⁸ K. Peters,⁸ T. Schroeder,⁸ M. Steinke,⁸ J. T. Boyd,⁹ J. P. Burke,⁹ N. Chevalier,⁹ W. N. Cottingham,⁹ M. P. Kelly,⁹ T. E. Latham,⁹ F. F. Wilson,⁹ T. Cuhadar-Donszelmann,¹⁰ C. Hearty,¹⁰ N. S. Knecht,¹⁰ T. S. Mattison,¹⁰ J. A. McKenna,¹⁰ D. Thiessen,¹⁰ A. Khan,¹¹ P. Kyberd,¹¹ L. Teodorescu,¹¹ A. E. Blinov,¹² V. E. Blinov,¹² V. P. Druzhinin,¹² V. B. Golubev,¹² V. N. Ivanchenko,¹² E. A. Kravchenko,¹² A. P. Onuchin,¹² S. I. Serednyakov,¹² Yu. I. Skovpen,¹² E. P. Solodov,¹² A. N. Yushkov,¹² D. Best,¹³ M. Bruinsma,¹³ M. Chao,¹³ I. Eschrich,¹³ D. Kirkby,¹³ A. J. Lankford,¹³ M. Mandelkern,¹³ R. K. Mommsen,¹³ W. Roethel,¹³ D. P. Stoker,¹³ C. Buchanan,¹⁴ B. L. Hartfel,¹⁴ A. J. R. Weinstein,¹⁴ S. D. Foulkes,¹⁵ J. W. Gary,¹⁵ O. Long,¹⁵ B. C. Shen,¹⁵ K. Wang,¹⁵ D. del Re,¹⁶ H. K. Hadavand,¹⁶ E. J. Hill,¹⁶ D. B. MacFarlane,¹⁶ H. P. Paar,¹⁶ Sh. Rahatlou,¹⁶ V. Sharma,¹⁶ J. W. Berryhill,¹⁷ C. Campagnari,¹⁷ A. Cunha,¹⁷ B. Dahmes,¹⁷ T. M. Hong,¹⁷ A. Lu,¹⁷ M. A. Mazur,¹⁷ J. D. Richman,¹⁷ W. Verkerke,¹⁷ T. W. Beck,¹⁸ A. M. Eisner,¹⁸ C. J. Flacco,¹⁸ C. A. Heusch,¹⁸ J. Kroseberg,¹⁸ W. S. Lockman,¹⁸ G. Nesom,¹⁸ T. Schalk,¹⁸ B. A. Schumm,¹⁸ A. Seiden,¹⁸ P. Spradlin,¹⁸ D. C. Williams,¹⁸ M. G. Wilson,¹⁸ J. Albert,¹⁹ E. Chen,¹⁹ G. P. Dubois-Felsmann,¹⁹ A. Dvoretzskii,¹⁹ D. G. Hitlin,¹⁹ I. Narsky,¹⁹ T. Piatenko,¹⁹ F. C. Porter,¹⁹ A. Ryd,¹⁹ A. Samuel,¹⁹ S. Yang,¹⁹ S. Jayatilke,²⁰ G. Mancinelli,²⁰ B. T. Meadows,²⁰ M. D. Sokoloff,²⁰ F. Blanc,²¹ P. Bloom,²¹ S. Chen,²¹ W. T. Ford,²¹ U. Nauenberg,²¹ A. Olivas,²¹ P. Rankin,²¹ W. O. Ruddick,²¹ J. G. Smith,²¹ K. A. Ulmer,²¹ J. Zhang,²¹ L. Zhang,²¹ A. Chen,²² E. A. Eckhart,²² J. L. Harton,²² A. Soffer,²² W. H. Toki,²² R. J. Wilson,²² Q. Zeng,²² B. Spaan,²³ D. Altenburg,²⁴ T. Brandt,²⁴ J. Brose,²⁴ M. Dickopp,²⁴ E. Feltresi,²⁴ A. Hauke,²⁴ H. M. Lacker,²⁴ E. Maly,²⁴ R. Nogowski,²⁴ S. Otto,²⁴ A. Petzold,²⁴ G. Schott,²⁴ J. Schubert,²⁴ K. R. Schubert,²⁴ R. Schwierz,²⁴ J. E. Sundermann,²⁴ D. Bernard,²⁵ G. R. Bonneaud,²⁵ P. Grenier,²⁵ S. Schrenk,²⁵ Ch. Thiebaux,²⁵ G. Vasileiadis,²⁵ M. Verderi,²⁵ D. J. Bard,²⁶ P. J. Clark,²⁶ F. Muheim,²⁶ S. Playfer,²⁶ Y. Xie,²⁶ M. Andreotti,²⁷ V. Azzolini,²⁷ D. Bettoni,²⁷ C. Bozzi,²⁷ R. Calabrese,²⁷ G. Cibinetto,²⁷ E. Luppi,²⁷ M. Negrini,²⁷ L. Piemontese,²⁷ A. Sarti,²⁷ F. Anulli,²⁸ R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,²⁸ M. Piccolo,²⁸ A. Zallo,²⁸ A. Buzzo,²⁹ R. Capra,²⁹ R. Contri,²⁹ G. Crosetti,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ C. Patrignani,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ S. Bailey,³⁰ G. Brandenburg,³⁰ K. S. Chaisanguanthum,³⁰ M. Morii,³⁰ E. Won,³⁰ R. S. Dubitzky,³¹ U. Langenegger,³¹ J. Marks,³¹ U. Uwer,³¹ W. Bhimji,³² D. A. Bowerman,³² P. D. Dauncey,³² U. Egede,³² J. R. Gaillard,³² G. W. Morton,³² J. A. Nash,³² M. B. Nikolich,³² G. P. Taylor,³² M. J. Charles,³³ G. J. Grenier,³³ U. Mallik,³³ A. K. Mohapatra,³³ J. Cochran,³⁴ H. B. Crawley,³⁴ J. Lamsa,³⁴ W. T. Meyer,³⁴ S. Prell,³⁴ E. I. Rosenberg,³⁴ A. E. Rubin,³⁴ J. Yi,³⁴ N. Arnaud,³⁵ M. Davier,³⁵ X. Giroux,³⁵ G. Grosdidier,³⁵ A. Höcker,³⁵ F. Le Diberder,³⁵ V. Lepeltier,³⁵ A. M. Lutz,³⁵ T. C. Petersen,³⁵ M. Pierini,³⁵ S. Plaszczyński,³⁵ M. H. Schune,³⁵ G. Wormser,³⁵ C. H. Cheng,³⁶ D. J. Lange,³⁶ M. C. Simani,³⁶ D. M. Wright,³⁶ A. J. Bevan,³⁷ C. A. Chavez,³⁷ J. P. Coleman,³⁷ I. J. Forster,³⁷ J. R. Fry,³⁷ E. Gabathuler,³⁷ R. Gamet,³⁷ D. E. Hutchcroft,³⁷ R. J. Parry,³⁷ D. J. Payne,³⁷ C. Touramanis,³⁷ C. M. Cormack,³⁸ F. Di Lodovico,³⁸ C. L. Brown,³⁹ G. Cowan,³⁹ R. L. Flack,³⁹ H. U. Flaecher,³⁹ M. G. Green,³⁹ P. S. Jackson,³⁹ T. R. McMahon,³⁹ S. Ricciardi,³⁹ F. Salvatore,³⁹ M. A. Winter,³⁹ D. Brown,⁴⁰ C. L. Davis,⁴⁰ J. Allison,⁴¹ N. R. Barlow,⁴¹ R. J. Barlow,⁴¹ M. C. Hodgkinson,⁴¹ G. D. Lafferty,⁴¹ M. T. Naisbit,⁴¹ J. C. Williams,⁴¹ C. Chen,⁴² A. Farbin,⁴² W. D. Hulsbergen,⁴² A. Jawahery,⁴² D. Kovalskyi,⁴² C. K. Lae,⁴² V. Lillard,⁴² D. A. Roberts,⁴² G. Blaylock,⁴³ C. Dallapiccola,⁴³ S. S. Hertzbach,⁴³ R. Kofler,⁴³

V. B. Koptchev,⁴³ T. B. Moore,⁴³ S. Saremi,⁴³ H. Staengle,⁴³ S. Willocq,⁴³ R. Cowan,⁴⁴ K. Koeneke,⁴⁴ G. Sciolla,⁴⁴ S. J. Sekula,⁴⁴ F. Taylor,⁴⁴ R. K. Yamamoto,⁴⁴ P. M. Patel,⁴⁵ S. H. Robertson,⁴⁵ A. Lazzaro,⁴⁶ V. Lombardo,⁴⁶ F. Palombo,⁴⁶ J. M. Bauer,⁴⁷ L. Cremaldi,⁴⁷ V. Eschenburg,⁴⁷ R. Godang,⁴⁷ R. Kroeger,⁴⁷ J. Reidy,⁴⁷ D. A. Sanders,⁴⁷ D. J. Summers,⁴⁷ H. W. Zhao,⁴⁷ S. Brunet,⁴⁸ D. Côté,⁴⁸ P. Taras,⁴⁸ H. Nicholson,⁴⁹ N. Cavallo,^{50,*} F. Fabozzi,^{50,*} C. Gatto,⁵⁰ L. Lista,⁵⁰ D. Monorchio,⁵⁰ P. Paolucci,⁵⁰ D. Piccolo,⁵⁰ C. Sciacca,⁵⁰ M. Baak,⁵¹ H. Bulten,⁵¹ G. Raven,⁵¹ H. L. Snoek,⁵¹ L. Wilden,⁵¹ C. P. Jessop,⁵² J. M. LoSecco,⁵² T. Allmendinger,⁵³ G. Benelli,⁵³ K. K. Gan,⁵³ K. Honscheid,⁵³ D. Hufnagel,⁵³ H. Kagan,⁵³ R. Kass,⁵³ T. Pulliam,⁵³ A. M. Rahimi,⁵³ R. Ter-Antonyan,⁵³ Q. K. Wong,⁵³ J. Brau,⁵⁴ R. Frey,⁵⁴ O. Igonkina,⁵⁴ M. Lu,⁵⁴ C. T. Potter,⁵⁴ N. B. Sinev,⁵⁴ D. Strom,⁵⁴ E. Torrence,⁵⁴ F. Colecchia,⁵⁵ A. Dorigo,⁵⁵ F. Galeazzi,⁵⁵ M. Margoni,⁵⁵ M. Morandin,⁵⁵ M. Posocco,⁵⁵ M. Rotondo,⁵⁵ F. Simonetto,⁵⁵ R. Stroili,⁵⁵ C. Voci,⁵⁵ M. Benayoun,⁵⁶ H. Briand,⁵⁶ J. Chauveau,⁵⁶ P. David,⁵⁶ L. Del Buono,⁵⁶ Ch. de la Vaissière,⁵⁶ O. Hamon,⁵⁶ M. J. J. John,⁵⁶ Ph. Leruste,⁵⁶ J. Malcès,⁵⁶ J. Ocariz,⁵⁶ L. Roos,⁵⁶ G. Therin,⁵⁶ P. K. Behera,⁵⁷ L. Gladney,⁵⁷ Q. H. Guo,⁵⁷ J. Panetta,⁵⁷ M. Biasini,⁵⁸ R. Covarelli,⁵⁸ M. Pioppi,⁵⁸ C. Angelini,⁵⁹ G. Batignani,⁵⁹ S. Bettarini,⁵⁹ M. Bondioli,⁵⁹ F. Bucci,⁵⁹ G. Calderini,⁵⁹ M. Carpinelli,⁵⁹ F. Forti,⁵⁹ M. A. Giorgi,⁵⁹ A. Lusiani,⁵⁹ G. Marchiori,⁵⁹ M. Morganti,⁵⁹ N. Neri,⁵⁹ E. Paoloni,⁵⁹ M. Rama,⁵⁹ G. Rizzo,⁵⁹ G. Simi,⁵⁹ J. Walsh,⁵⁹ M. Haire,⁶⁰ D. Judd,⁶⁰ K. Paick,⁶⁰ D. E. Wagoner,⁶⁰ N. Danielson,⁶¹ P. Elmer,⁶¹ Y. P. Lau,⁶¹ C. Lu,⁶¹ V. Miftakov,⁶¹ J. Olsen,⁶¹ A. J. S. Smith,⁶¹ A. V. Telnov,⁶¹ F. Bellini,⁶² G. Cavoto,^{61,62} A. D'Orazio,⁶² E. Di Marco,⁶² R. Faccini,⁶² F. Ferrarotto,⁶² F. Ferroni,⁶² M. Gaspero,⁶² L. Li Gioi,⁶² M. A. Mazzone,⁶² S. Morganti,⁶² G. Piredda,⁶² F. Polci,⁶² F. Safai Tehrani,⁶² C. Voena,⁶² S. Christ,⁶³ H. Schröder,⁶³ G. Wagner,⁶³ R. Waldi,⁶³ T. Adye,⁶⁴ N. De Groot,⁶⁴ B. Franek,⁶⁴ G. P. Gopal,⁶⁴ E. O. Olaiya,⁶⁴ R. Aleksan,⁶⁵ S. Emery,⁶⁵ A. Gaidot,⁶⁵ S. F. Ganzhur,⁶⁵ P.-F. Giraud,⁶⁵ G. Graziani,⁶⁵ G. Hamel de Monchenault,⁶⁵ W. Kozanecki,⁶⁵ M. Legendre,⁶⁵ G. W. London,⁶⁵ B. Mayer,⁶⁵ G. Vasseur,⁶⁵ Ch. Yèche,⁶⁵ M. Zito,⁶⁵ M. V. Purohit,⁶⁶ A. W. Weidemann,⁶⁶ J. R. Wilson,⁶⁶ F. X. Yumiceva,⁶⁶ T. Abe,⁶⁷ M. T. Allen,⁶⁷ D. Aston,⁶⁷ R. Bartoldus,⁶⁷ N. Berger,⁶⁷ A. M. Boyarski,⁶⁷ O. L. Buchmueller,⁶⁷ R. Claus,⁶⁷ M. R. Convery,⁶⁷ M. Cristinziani,⁶⁷ G. De Nardo,⁶⁷ J. C. Dingfelder,⁶⁷ D. Dong,⁶⁷ J. Dorfan,⁶⁷ D. Dujmic,⁶⁷ W. Dunwoodie,⁶⁷ S. Fan,⁶⁷ R. C. Field,⁶⁷ T. Glanzman,⁶⁷ S. J. Gowdy,⁶⁷ T. Hadig,⁶⁷ V. Halyo,⁶⁷ C. Hast,⁶⁷ T. Hryn'ova,⁶⁷ W. R. Innes,⁶⁷ M. H. Kelsey,⁶⁷ P. Kim,⁶⁷ M. L. Kocian,⁶⁷ D. W. G. S. Leith,⁶⁷ J. Libby,⁶⁷ S. Luitz,⁶⁷ V. Luth,⁶⁷ H. L. Lynch,⁶⁷ H. Marsiske,⁶⁷ R. Messner,⁶⁷ D. R. Muller,⁶⁷ C. P. O'Grady,⁶⁷ V. E. Ozcan,⁶⁷ A. Perazzo,⁶⁷ M. Perl,⁶⁷ B. N. Ratcliff,⁶⁷ A. Roodman,⁶⁷ A. A. Salnikov,⁶⁷ R. H. Schindler,⁶⁷ J. Schwiening,⁶⁷ A. Snyder,⁶⁷ A. Soha,⁶⁷ J. Stelzer,⁶⁷ J. Strube,^{54,67} D. Su,⁶⁷ M. K. Sullivan,⁶⁷ J. M. Thompson,⁶⁷ J. Va'vra,⁶⁷ S. R. Wagner,⁶⁷ M. Weaver,⁶⁷ W. J. Wisniewski,⁶⁷ M. Wittgen,⁶⁷ D. H. Wright,⁶⁷ A. K. Yarritu,⁶⁷ C. C. Young,⁶⁷ P. R. Burchat,⁶⁸ A. J. Edwards,⁶⁸ S. A. Majewski,⁶⁸ B. A. Petersen,⁶⁸ C. Roat,⁶⁸ M. Ahmed,⁶⁹ S. Ahmed,⁶⁹ M. S. Alam,⁶⁹ J. A. Ernst,⁶⁹ M. A. Saeed,⁶⁹ M. Saleem,⁶⁹ F. R. Wappler,⁶⁹ W. Bugg,⁷⁰ M. Krishnamurthy,⁷⁰ S. M. Spanier,⁷⁰ R. Eckmann,⁷¹ H. Kim,⁷¹ J. L. Ritchie,⁷¹ A. Satpathy,⁷¹ R. F. Schwitters,⁷¹ J. M. Izen,⁷² I. Kitayama,⁷² X. C. Lou,⁷² S. Ye,⁷² F. Bianchi,⁷³ M. Bona,⁷³ F. Gallo,⁷³ D. Gamba,⁷³ L. Bosisio,⁷⁴ C. Cartaro,⁷⁴ F. Cossutti,⁷⁴ G. Della Ricca,⁷⁴ S. Dittongo,⁷⁴ S. Grancagnolo,⁷⁴ L. Lanceri,⁷⁴ P. Poropat,^{74,†} L. Vitale,⁷⁴ G. Vuagnin,⁷⁴ F. Martinez-Vidal,^{2,75} R. S. Panvini,^{76,†} Sw. Banerjee,⁷⁷ B. Bhuyan,⁷⁷ C. M. Brown,⁷⁷ D. Fortin,⁷⁷ K. Hamano,⁷⁷ P. D. Jackson,⁷⁷ R. Kowalewski,⁷⁷ J. M. Roney,⁷⁷ R. J. Sobie,⁷⁷ J. J. Back,⁷⁸ P. F. Harrison,⁷⁸ G. B. Mohanty,⁷⁸ H. R. Band,⁷⁹ X. Chen,⁷⁹ B. Cheng,⁷⁹ S. Dasu,⁷⁹ M. Datta,⁷⁹ A. M. Eichenbaum,⁷⁹ K. T. Flood,⁷⁹ M. Graham,⁷⁹ J. J. Hollar,⁷⁹ J. R. Johnson,⁷⁹ P. E. Kutter,⁷⁹ H. Li,⁷⁹ R. Liu,⁷⁹ A. Mihalyi,⁷⁹ Y. Pan,⁷⁹ R. Prepost,⁷⁹ P. Tan,⁷⁹ J. H. von Wimmersperg-Toeller,⁷⁹ J. Wu,⁷⁹ S. L. Wu,⁷⁹ Z. Yu,⁷⁹ M. G. Greene,⁸⁰ and H. Neal⁸⁰

(The BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France

²IFAE, Universitat Autònoma de Barcelona, E-08193 Bellaterra, Barcelona, Spain

³Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy

⁴Institute of High Energy Physics, Beijing 100039, China

⁵University of Bergen, Inst. of Physics, N-5007 Bergen, Norway

⁶Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720, USA

⁷University of Birmingham, Birmingham, B15 2TT, United Kingdom

⁸Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany

⁹University of Bristol, Bristol BS8 1TL, United Kingdom

¹⁰University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1

¹¹Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom

¹²Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia

¹³University of California at Irvine, Irvine, California 92697, USA

- ¹⁴ *University of California at Los Angeles, Los Angeles, California 90024, USA*
- ¹⁵ *University of California at Riverside, Riverside, California 92521, USA*
- ¹⁶ *University of California at San Diego, La Jolla, California 92093, USA*
- ¹⁷ *University of California at Santa Barbara, Santa Barbara, California 93106, USA*
- ¹⁸ *University of California at Santa Cruz, Institute for Particle Physics, Santa Cruz, California 95064, USA*
- ¹⁹ *California Institute of Technology, Pasadena, California 91125, USA*
- ²⁰ *University of Cincinnati, Cincinnati, Ohio 45221, USA*
- ²¹ *University of Colorado, Boulder, Colorado 80309, USA*
- ²² *Colorado State University, Fort Collins, Colorado 80523, USA*
- ²³ *Universität Dortmund, Institut für Physik, D-44221 Dortmund, Germany*
- ²⁴ *Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany*
- ²⁵ *Ecole Polytechnique, LLR, F-91128 Palaiseau, France*
- ²⁶ *University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom*
- ²⁷ *Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy*
- ²⁸ *Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy*
- ²⁹ *Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy*
- ³⁰ *Harvard University, Cambridge, Massachusetts 02138, USA*
- ³¹ *Universität Heidelberg, Physikalisches Institut, Philosophenweg 12, D-69120 Heidelberg, Germany*
- ³² *Imperial College London, London, SW7 2AZ, United Kingdom*
- ³³ *University of Iowa, Iowa City, Iowa 52242, USA*
- ³⁴ *Iowa State University, Ames, Iowa 50011-3160, USA*
- ³⁵ *Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France*
- ³⁶ *Lawrence Livermore National Laboratory, Livermore, California 94550, USA*
- ³⁷ *University of Liverpool, Liverpool L69 7ZE, United Kingdom*
- ³⁸ *Queen Mary, University of London, E1 4NS, United Kingdom*
- ³⁹ *University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom*
- ⁴⁰ *University of Louisville, Louisville, Kentucky 40292, USA*
- ⁴¹ *University of Manchester, Manchester M13 9PL, United Kingdom*
- ⁴² *University of Maryland, College Park, Maryland 20742, USA*
- ⁴³ *University of Massachusetts, Amherst, Massachusetts 01003, USA*
- ⁴⁴ *Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139, USA*
- ⁴⁵ *McGill University, Montréal, Quebec, Canada H3A 2T8*
- ⁴⁶ *Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy*
- ⁴⁷ *University of Mississippi, University, Mississippi 38677, USA*
- ⁴⁸ *Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Quebec, Canada H3C 3J7*
- ⁴⁹ *Mount Holyoke College, South Hadley, Massachusetts 01075, USA*
- ⁵⁰ *Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126, Napoli, Italy*
- ⁵¹ *NIKHEF, National Institute for Nuclear Physics and High Energy Physics, NL-1009 DB Amsterdam, The Netherlands*
- ⁵² *University of Notre Dame, Notre Dame, Indiana 46556, USA*
- ⁵³ *Ohio State University, Columbus, Ohio 43210, USA*
- ⁵⁴ *University of Oregon, Eugene, Oregon 97403, USA*
- ⁵⁵ *Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy*
- ⁵⁶ *Universités Paris VI et VII, Laboratoire de Physique Nucléaire et de Hautes Energies, F-75252 Paris, France*
- ⁵⁷ *University of Pennsylvania, Philadelphia, Pennsylvania 19104, USA*
- ⁵⁸ *Università di Perugia, Dipartimento di Fisica and INFN, I-06100 Perugia, Italy*
- ⁵⁹ *Università di Pisa, Dipartimento di Fisica, Scuola Normale Superiore and INFN, I-56127 Pisa, Italy*
- ⁶⁰ *Prairie View A&M University, Prairie View, Texas 77446, USA*
- ⁶¹ *Princeton University, Princeton, New Jersey 08544, USA*
- ⁶² *Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy*
- ⁶³ *Universität Rostock, D-18051 Rostock, Germany*
- ⁶⁴ *Rutherford Appleton Laboratory, Chilton, Didcot, Oxon, OX11 0QX, United Kingdom*
- ⁶⁵ *DSM/Dapnia, CEA/Saclay, F-91191 Gif-sur-Yvette, France*
- ⁶⁶ *University of South Carolina, Columbia, South Carolina 29208, USA*
- ⁶⁷ *Stanford Linear Accelerator Center, Stanford, California 94309, USA*
- ⁶⁸ *Stanford University, Stanford, California 94305-4060, USA*
- ⁶⁹ *State University of New York, Albany, New York 12222, USA*
- ⁷⁰ *University of Tennessee, Knoxville, Tennessee 37996, USA*
- ⁷¹ *University of Texas at Austin, Austin, Texas 78712, USA*
- ⁷² *University of Texas at Dallas, Richardson, Texas 75083, USA*
- ⁷³ *Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy*
- ⁷⁴ *Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy*
- ⁷⁵ *IFIC, Universitat de Valencia-CSIC, E-46071 Valencia, Spain*
- ⁷⁶ *Vanderbilt University, Nashville, Tennessee 37235, USA*
- ⁷⁷ *University of Victoria, Victoria, British Columbia, Canada V8W 3P6*

⁷⁸*Department of Physics, University of Warwick, Coventry CV4 7AL, United Kingdom*

⁷⁹*University of Wisconsin, Madison, Wisconsin 53706, USA*

⁸⁰*Yale University, New Haven, Connecticut 06511, USA*

(Dated: February 7, 2008)

We measure the time-dependent CP asymmetry parameters in $B^0 \rightarrow K^+K^-K^0$ based on a data sample of approximately 227 million B -meson pairs recorded at the $\Upsilon(4S)$ resonance with the BABAR detector at the PEP-II B -meson Factory at SLAC. We reconstruct two-body B^0 decays to $\phi(1020)K_S^0$ and $\phi(1020)K_L^0$, and the three-body decay $K^+K^-K_S^0$ with $\phi(1020)K_S^0$ excluded. For the $B^0 \rightarrow \phi K^0$ decays, we measure $\sin 2\beta_{\text{eff}}(\phi K^0) = +0.50 \pm 0.25$ (stat) $^{+0.07}_{-0.04}$ (syst). The $B^0 \rightarrow K^+K^-K_S^0$ decays are dominated by K^+K^- S -wave, as determined from an angular analysis; we measure $\sin 2\beta_{\text{eff}}(K^+K^-K_S^0) = +0.55 \pm 0.22$ (stat) ± 0.04 (syst) ± 0.11 (CP), where the last error is due to the uncertainty in the fraction of CP -even contributions to the decay amplitude. We find no evidence for direct CP violation.

PACS numbers: 13.25.Hw, 12.15.Hh, 11.30.Er

In the Standard Model (SM) of particle physics, the decays $B^0 \rightarrow K^+K^-K^0$ [1] are dominated by $b \rightarrow s\bar{s}s$ gluonic penguin amplitudes, but can also be affected by amplitudes that are suppressed by elements of the Cabibbo–Kobayashi–Maskawa (CKM) quark mixing matrix [2]. These CKM-suppressed amplitudes cannot be precisely known in a model-independent way [3], but are in general expected to be small [4]. Let $2\beta_{\text{eff}}$ be the CP -violating phase difference between decays with and without mixing, and $\beta = \arg(-V_{cd}V_{cb}^*/V_{td}V_{tb}^*)$ where V_{ij} are elements of the CKM quark mixing matrix. The difference $|\beta - \beta_{\text{eff}}|$ is expected to be nearly zero, with theoretical uncertainties of a few degrees for $B^0 \rightarrow \phi K^0$ [5]. Larger uncertainties exist for $B^0 \rightarrow K^+K^-K_S^0$ with $B^0 \rightarrow \phi K_S^0$ decays excluded, due in part to an extra CKM-suppressed tree amplitude contribution [4].

Since additional decay diagrams with non-SM particles and interactions introducing new CP -violating phases may contribute to β_{eff} , measurements of $\sin 2\beta_{\text{eff}}$ in these channels and their comparisons with the SM expectation are sensitive probes for physics beyond the SM [4]. The value of $\sin 2\beta$ has been measured in $B^0 \rightarrow J/\psi K_S^0$ [6, 7] with an average of 0.742 ± 0.037 . The BABAR and Belle collaborations have measured $\sin 2\beta_{\text{eff}}$ in ϕK^0 ($+0.47 \pm 0.34^{+0.08}_{-0.06}$ with 114 million $B\bar{B}$ pairs [8] and $-0.96 \pm 0.50^{+0.09}_{-0.11}$ with 152 million $B\bar{B}$ pairs (ϕK_S^0 only) [9], respectively), and in $K^+K^-K_S^0$ excluding ϕK_S^0 ($+0.57 \pm 0.26 \pm 0.04^{+0.17}_{-0.00}$ with 122 million $B\bar{B}$ pairs [10] and $+0.51 \pm 0.26 \pm 0.05^{+0.18}_{-0.00}$ with 152 million $B\bar{B}$ pairs [9], respectively).

At B factories, the neutral B mesons are exclusively produced in pairs. We select events for which one B (B_{rec}) is reconstructed as $B^0 \rightarrow K^+K^-K^0$ and the other (B_{tag}) is partially reconstructed as either B^0 or \bar{B}^0 . We define $\Delta t = t_{\text{rec}} - t_{\text{tag}}$ to be the difference between the proper decay times of the B mesons. The decay rate $f_+(f_-)$ for the final state f when the B_{tag} decays as a B^0

(\bar{B}^0) is given by

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau_{B^0}}}{4\tau_{B^0}} [1 \pm S_f \sin(\Delta m_d \Delta t) \mp C_f \cos(\Delta m_d \Delta t)], \quad (1)$$

where τ_{B^0} is the B^0 lifetime and Δm_d is the B^0 – \bar{B}^0 mixing frequency. The parameter S_f is non-zero if there is CP violation in the interference between decays with and without mixing, while a non-zero value for C_f would entail direct CP violation. In the limit where the CKM-suppressed amplitudes do not contribute, the SM predicts no direct CP violation ($C_f = 0$) since the dominant decay amplitudes have the same CP -violating phase, and that $S_f = -\eta_f \times \sin 2\beta_{\text{eff}}$. For $B^0 \rightarrow \phi K_S^0$ decays, the effective eigenvalue $\eta_f = -1$, for $B^0 \rightarrow \phi K_L^0$ $\eta_f = +1$. For $B^0 \rightarrow K^+K^-K_S^0$ decays, $\eta_f = 2f_{\text{even}} - 1$, where f_{even} is the fraction of CP -even contributions to the $B^0 \rightarrow K^+K^-K_S^0$ amplitude. Then the value of η_f depends on the angular momentum of the K^+K^- system: it is -1 for relative P -wave and +1 for S -wave.

In this paper, we present a measurement of $\sin 2\beta_{\text{eff}}$ with almost twice the number of events as for the previous BABAR results [8, 10]. We reconstruct B^0 candidates in two independent modes, ϕK^0 (with the K^0 either a K_L^0 or a K_S^0) and $K^+K^-K_S^0$ (with the ϕ mass region excluded). K_S^0 's are detected via their $\pi^+\pi^-$ decay only. We extract the CP asymmetry parameters using extended maximum-likelihood fits. Using an angular moment analysis [11], we extract the K^+K^- P -wave fractions in the data. These fractions are used to check the assumption that $\eta_f = -1$ for ϕK_S^0 and +1 for ϕK_L^0 by bounding the S -wave contamination in the ϕ mass region, and to measure η_f for $K^+K^-K_S^0$.

This analysis is based on 227 million $B\bar{B}$ pairs collected with the BABAR detector [12] at the PEP-II asymmetric-energy e^+e^- storage rings at SLAC, operating at the $\Upsilon(4S)$ resonance (center-of-mass (c.m.) energy $\sqrt{s} = 10.58$ GeV). In Ref. 12 we describe the silicon vertex tracker (SVT) and drift chamber (DCH) used for track and vertex reconstruction, the electromagnetic calorime-

ter (EMC) and instrumented flux return (IFR) used for K_L^0 reconstruction, and the detector of internally reflected Cherenkov light (DIRC), which, together with the EMC, the IFR, and the ionization dE/dx from the SVT and DCH, is used for particle identification.

The B^0 -candidate reconstruction and selection is similar to that described in Refs. 8 and 10. We consider a K^+K^- pair to be a ϕ candidate if its invariant mass is within $15 \text{ MeV}/c^2$ (about three times the apparent width in the K^+K^- invariant mass spectrum) of the central ϕ mass value [13]. For a given $B^0\bar{B}^0$ meson pair, we obtain Δt from the measured distance between the fully reconstructed B_{rec} meson decay point and the B_{tag} decay point along the beam direction, and the known boost of the $\Upsilon(4S)$ system ($\beta\gamma=0.56$). A multivariate tagging algorithm determines the flavor of the B_{tag} meson [6] and classifies it in one of seven mutually exclusive tagging categories.

We use two kinematic variables to discriminate between signal B decays and combinatorial background. The energy difference between the measured e^+e^- c.m. energy of the B candidate and $\sqrt{s}/2$ is ΔE . Its distribution peaks at zero for signal, with a width of about 20 MeV for ϕK_S^0 and $K^+K^-K_S^0$. The width is only about 3 MeV for ϕK_L^0 , because for this mode we constrain the B^0 candidate's mass to the nominal value [13]. The beam-energy-substituted mass, m_{ES} , is used for candidates without a K_L^0 . It is defined as $m_{\text{ES}} \equiv \sqrt{(s/2 + \mathbf{p}_i \cdot \mathbf{p}_B)^2/E_i^2 - \mathbf{p}_B^2}$, where the B momentum \mathbf{p}_B and the four-momentum of the initial state (E_i, \mathbf{p}_i) are defined in the laboratory frame. It peaks at the B^0 mass for signal, with a width of about 3 MeV . For ϕK_S^0 candidates, we require $|\Delta E| < 100 \text{ MeV}$ and $m_{\text{ES}} > 5.21 \text{ GeV}/c^2$; for ϕK_L^0 candidates, we require $|\Delta E| < 80 \text{ MeV}$; and for $K^+K^-K_S^0$ candidates, we require $|\Delta E| < 200 \text{ MeV}$ and $m_{\text{ES}} > 5.2 \text{ GeV}/c^2$.

The dominant background is continuum $e^+e^- \rightarrow q\bar{q}$ ($q = u, d, s, c$) events; these tend to be jet-like in the e^+e^- c.m. frame, while B decays tend to be spherical. To enhance discrimination between signal and continuum, we use Fisher discriminants (\mathcal{F}) to combine four event-shape-related variables [8, 10]. The other background originates from B decays. For the ϕK^0 final state, opposite- CP contributions from the $K^+K^-K^0$ final state (K^+K^- S -wave) are estimated from data with a moment analysis [11] (see below) to be less than 6.6% at 95% confidence level. The mode ϕK_L^0 has additional background. Its dominant CP contamination is from the mode $\phi K^{*0} \rightarrow \phi K_L^0 \pi^0$, for which we expect approximately eight events in the region $|\Delta E| < 10 \text{ MeV}$. In the final likelihood fit we explicitly parameterize backgrounds from B decays both with and without charm.

For the $K^+K^-K_S^0$ mode, we apply invariant mass cuts to suppress background from B decays that proceed through a $b \rightarrow c$ transition, namely those containing D^0 , J/ψ , χ_{c0} , or $\psi(2S)$ decaying into K^+K^- , or D^+ or D_s^+

decaying into $K^+K_S^0$. Finally, to suppress B decays into final states with pions, we require the rate for a charged pion to be misidentified as a kaon to be less than 2% .

A total of 4,300, 8,238, and 27,368 events have a candidate that passes the ϕK_S^0 , ϕK_L^0 , or $K^+K^-K_S^0$ selection criteria, respectively. From simulation, we find the final selection efficiencies for signal to be 40% , 20% and 26% , respectively.

We extract the $K^+K^-K_S^0$, ϕK_S^0 , and ϕK_L^0 event yields and CP parameters with two extended maximum-likelihood fits. One is to the $K^+K^-K_S^0$ candidates; the other is to both the ϕK_S^0 and ϕK_L^0 candidates, with the assumption that $C_{\phi K_S^0} = C_{\phi K_L^0}$ and $S_{\phi K_S^0} = -S_{\phi K_L^0}$. We verified the fit procedure for the ϕK^0 mode with samples of ϕK^+ and $J/\psi K^0$ events. We found for the former a null asymmetry as expected, and for the latter results that are consistent with previous measurements [6]. We verified the fit procedure for the $K^+K^-K_S^0$ mode with a sample of $K_S^0 K_S^0 K^+$ events, for which we found a null asymmetry as expected.

The likelihood function used in each extended maximum-likelihood fit to its N_k candidates tagged in category k is

$$\mathcal{L}_k = e^{-N'_k} \prod_{i=1}^{N_k} \left\{ N_S \epsilon_k \mathcal{P}_{i,k}^S + N_{C,k} \mathcal{P}_{i,k}^C + \sum_{j=1}^{n_B} N_{B,j} \epsilon_{j,k} \mathcal{P}_{i,j,k}^B \right\} \quad (2)$$

where N'_k is the sum of the signal, continuum, and n_B B -background yields tagged in category k ; N_S is the number of ϕK_S^0 , ϕK_L^0 , or $K^+K^-K_S^0$ signal events; ϵ_k is the fraction of signal events tagged in category k ; $N_{C,k}$ is the number of continuum background events tagged in category k ; $N_{B,j}$ is the number of B -background events of class j ; and $\epsilon_{j,k}$ is the fraction of B -background events of class j tagged in category k . Each B -background event comprises similar B decays. The B -background event yields are fixed parameters and are zero for the ϕK_S^0 sample. The total likelihood \mathcal{L} is the product of the likelihoods for each tagging category.

The probability density functions (PDFs) \mathcal{P}_k^S , \mathcal{P}_k^C , and $\mathcal{P}_{j,k}^B$, for signal, continuum background, and B -background class j , respectively, are the products of the PDFs of the discriminating variables. The signal PDF is thus given for the $K^+K^-K_S^0$ sample by $\mathcal{P}(m_{\text{ES}}) \cdot \mathcal{P}(\Delta E) \cdot \mathcal{P}(\mathcal{F}) \cdot \mathcal{P}(\Delta t; \sigma_{\Delta t})$, for the ϕK_S^0 sample by $\mathcal{P}(m_{\text{ES}}) \cdot \mathcal{P}(\Delta E) \cdot \mathcal{P}(\mathcal{F}) \cdot \mathcal{P}(m_{KK}) \cdot \mathcal{P}(\cos \theta_H) \cdot \mathcal{P}(\Delta t; \sigma_{\Delta t})$, and for the ϕK_L^0 sample by $\mathcal{P}(\Delta E) \cdot \mathcal{P}(\mathcal{F}) \cdot \mathcal{P}(m_{KK}) \cdot \mathcal{P}(\cos \theta_H) \cdot \mathcal{P}(\Delta t; \sigma_{\Delta t})$, where θ_H is the angle between the K^+ candidate and the parent B_{rec} flight direction in the K^+K^- rest frame. The quantity $\sigma_{\Delta t}$ is the uncertainty in the measurement of Δt for a given event. The time-dependent CP parameters defined in Eq. (1), diluted by the effects of mistagging and the Δt resolution, are contained in $\mathcal{P}_k^S(\Delta t, \sigma_{\Delta t})$. As in our $J/\psi K_S^0$ analysis [6], the Δt -resolution function for signal and B -background

events is a sum of three Gaussian distributions, which have two distinct means as well as three distinct widths. The widths are the error of the measured Δt scaled by three independent factors.

In the fits to data, we leave unconstrained the parameters describing the CP asymmetry, the Δt -resolution functions, the tagging characteristics, and the event yields. We also leave unconstrained the means of the signal m_{ES} and ΔE Gaussian PDFs, the widths of the signal ΔE PDFs, the mean of the signal m_{KK} PDF (which is parameterized by a relativistic P -wave Breit-Wigner function), and all parameters of the $K^+K^-K_s^0$ candidates' signal PDF for \mathcal{F} . We take from simulation any other parameters of the m_{ES} , ΔE , \mathcal{F} , $\cos\theta_H$, and m_{KK} PDFs for signal and B background. The parameters describing the signal and B -background Δt -resolution function are determined by a simultaneous fit to an independent sample of reconstructed B^0 decays to flavor eigenstates, with more than 100,000 events [6]. We use the world-averaged values for τ_{B^0} and Δm_d [13]. The fits to the ϕK^0 and $K^+K^-K_s^0$ candidates have a total of 35 and 34 free parameters, respectively.

We use an angular moment analysis based on the $\cos\theta_H$ distribution to extract the $K^+K^-K_s^0$ CP content, and also to bound the S -wave contamination in the ϕ mass region. In this approach, we expand the decay distribution for a given K^+K^- invariant mass in terms of moments $\langle P_\ell \rangle$ of conveniently normalized Legendre polynomials $P_\ell(\cos\theta_H)$:

$$|\mathcal{A}(m_{KK})|^2 = \sum_\ell \langle P_\ell \rangle \cdot P_\ell(\cos\theta_H), \quad (3)$$

where $\mathcal{A}(m_{KK})$ is the mass-dependent decay amplitude. We normalize $P_\ell(\cos\theta_H)$ such that the integral of $P_\ell(\cos\theta_H)^2$ over $\cos\theta_H$ from -1 to 1 equals unity. We extract the moments by summing over all events:

$$\langle P_\ell \rangle = \sum_j P_\ell(\cos\theta_{H,j}) \mathcal{W}_j / \varepsilon_j, \quad (4)$$

where \mathcal{W}_j is the weight for event j to belong to the signal decay and is calculated by the sPlot technique of Ref. 14. The efficiency ε_j is evaluated from a large MC sample in bins of m_{KK} and $\cos\theta_H$. Limiting ourselves to the two lowest partial waves, we can write the total decay amplitude in terms of the S -wave (CP -even) and the P -wave (CP -odd) amplitudes,

$$\mathcal{A}(m_{KK}) \approx \mathcal{A}_S(m_{KK})P_0(\cos\theta_H) + e^{i\phi_p} \mathcal{A}_P(m_{KK})P_1(\cos\theta_H), \quad (5)$$

where ϕ_p is the relative phase between the real partial-wave amplitudes $\mathcal{A}_S(m_{KK})$ and $\mathcal{A}_P(m_{KK})$. If we compare Eq. (5) to Eq. (3), we can relate the moments (of order $\ell \leq 2$) to the wave intensities and thus to the total

fraction of CP -even events, f_{even} , as

$$\begin{aligned} f_{\text{even}} &= \frac{\mathcal{A}_S(m_{KK})^2}{\mathcal{A}_S(m_{KK})^2 + \mathcal{A}_P(m_{KK})^2} \\ &= 1 - \sqrt{\frac{5}{4}} \frac{\langle P_2 \rangle}{\langle P_0 \rangle}, \end{aligned} \quad (6)$$

where $\mathcal{A}_S(m_{KK})^2$ and $\mathcal{A}_P(m_{KK})^2$ are the S - and P -wave intensities, respectively. In the normalization, the total number of signal events is $\sqrt{2} \langle P_0 \rangle$.

TABLE I: Systematic uncertainties on the CP parameters.

Source	$S_{\phi K}$	$C_{\phi K}$	S_{KKK}	C_{KKK}
Detector effects	± 0.02	± 0.02	± 0.02	± 0.01
DCSD	± 0.01	± 0.03	± 0.00	± 0.03
Fit bias	± 0.01	± 0.01	± 0.02	± 0.01
$B^0\text{-}\bar{B}^0$ tagging	± 0.01	± 0.02	± 0.00	± 0.01
S -wave contamination	$+0.06$	± 0.02	-	-
Other	± 0.03	± 0.02	± 0.01	± 0.01
Total	$^{+0.07}_{-0.04}$	± 0.05	± 0.03	± 0.04

Systematic errors on the CP -asymmetry parameters are listed in Table I. We account for uncertainties in the Δt resolution, the beam-spot position, and the detector alignment. We also estimate errors due to the effect of doubly CKM-suppressed decays (DCSD) of the B_{tag} [15]. The uncertainty due to possible biases in the fit procedure is conservative and includes effects on the CP parameters of correlations among the fit variables, which have been determined with full-detector MC simulations. Uncertainties in the $B^0\text{-}\bar{B}^0$ tagging efficiency in both signal and background are also included. Finally, we account for errors due to the CP content of the background, uncertainties in the PDF parameterization, and the uncertainties of τ_{B^0} and Δm_d [13]. For each mode we add the individual contributions in quadrature to obtain the total systematic uncertainty.

We also consider the systematic error due to the CP -even fraction of the $K^+K^-K_s^0$ mode. We do not find evidence for the existence of higher moments $\langle P_\ell \rangle$, $\ell = 3 \dots 6$, which could arise from intermediate D -wave decays into K^+K^- or decays proceeding through an isospin-1 resonance into $K^\pm K_s^0$. Nevertheless, we estimate a systematic error from the D -wave by examining the $\langle P_2 \rangle$ moment in the K^+K^- mass region (1.1–1.7) GeV/ c^2 , corresponding to the $f_2(1270)$, $a_2(1320)^0$, and $f_2'(1525)$ resonances, and assuming that $\langle P_2 \rangle$ arises only from D -wave and S - D interference. Since the moment itself is consistent with zero, we assign a systematic error of 4% based on the $\langle P_2 \rangle$ error. We account for the possible presence of $a_0(980)^+$, $a_0(1450)^+$, and $a_2(1320)^+$ in the $K^\pm K_s^0$ subsystem (4.6%). We also estimate a bias due to the modeling of the efficiency from MC events (2.5%). We find the total systematic error on f_{even} to be ± 0.06 . This leads to a systematic error on $\sin 2\beta_{\text{eff}}$ of ± 0.11 .

Table II shows the measured CP parameters and yields from the final extended maximum-likelihood fits. Note that when fitting $\sin 2\beta_{\text{eff}}$ for $K^+K^-K_s^0$, we constrain C_{KKK} to zero. All yields are consistent with our previously measured branching fractions [10, 16]. Figure 1 shows the signal-enhanced distributions of m_{ES} for ϕK_s^0 and $K^+K^-K_s^0$ events and of ΔE for ϕK_L^0 events, together with the result from the final extended maximum-likelihood fits. Figure 2 shows the time-dependent asymmetry distributions. As a cross check, we also fit ϕK_s^0 and ϕK_L^0 separately. Our fit to only ϕK_s^0 events gives $S = 0.29 \pm 0.31$ and $C = -0.07 \pm 0.27$. Our fit to only ϕK_L^0 events gives $S = 1.05 \pm 0.51$ and $C = 0.31 \pm 0.49$.




FIG. 1: Distributions of (a) m_{ES} for ϕK_s^0 candidates, (b) ΔE for ϕK_L^0 candidates, and (c) m_{ES} for $K^+K^-K_s^0$ candidates excluding ϕK_s^0 , together with the results from the final extended maximum-likelihood fits after applying a requirement on the ratio of signal likelihood to signal-plus-background likelihood (computed without the displayed variable) to reduce the background. The requirement is chosen to roughly maximize $N_S^2/(N_S + N_C)$ where N_C is the total number of continuum events, and is applied only for the purpose of making these plots. The curves are projections from the likelihood fits for total yield (solid lines), continuum background (short dashed lines), and total background (long dashes in (b) only). The efficiency of the likelihood-ratio cut is (a) 79% for signal and 5% for background, (b) 35% for signal, 16% for B -background, and 3% for continuum background, and (c) 77% for signal and 5% for background.

For the $K^+K^-K_s^0$ final state including the ϕ mass region, the distributions of the S - and P -wave intensities, and the CP -even fraction, as a function of K^+K^- invariant mass, are shown in Fig. 3. The total fraction of CP -even events with the ϕ mass region excluded is

TABLE II: CP -asymmetry parameters and yields from the final extended maximum-likelihood fits, as well as the fraction of CP -even contributions to the amplitude, f_{even} , which is assumed to be zero for ϕK_s^0 and unity ϕK_L^0 . The first errors are statistical, and the second are systematic; the third error on $\sin 2\beta_{\text{eff}}$ for $K^+K^-K_s^0$ is due to the uncertainty in the CP content. The values of S and C are fit simultaneously for the ϕK_s^0 and ϕK_L^0 candidates; the sign of S for ϕK_s^0 is shown. When finding $\sin 2\beta_{\text{eff}}$ for $K^+K^-K_s^0$, we constrain C_{KKK} to 0.

	ϕK^0		$K^+K^-K_s^0$
	ϕK_s^0	ϕK_L^0	(no ϕK_s^0)
$\sin 2\beta_{\text{eff}}$	$+0.50 \pm 0.25^{+0.07}_{-0.04}$		$+0.55 \pm 0.22 \pm 0.04 \pm 0.11$
f_{even}	0	1	$0.89 \pm 0.08 \pm 0.06$
S	$+0.50 \pm 0.25^{+0.07}_{-0.04}$		$-0.42 \pm 0.17 \pm 0.03$
C	$0.00 \pm 0.23 \pm 0.05$		$+0.10 \pm 0.14 \pm 0.04$
Yield	114 ± 12	98 ± 18	452 ± 28




FIG. 2: The time-dependent asymmetry distributions for (a) ϕK_s^0 , (b) ϕK_L^0 , and (c) $K^+K^-K_s^0$ with no ϕK_s^0 decays. The asymmetry is defined as $A_{B^0/\bar{B}^0} = (N_{B^0} - N_{\bar{B}^0}) / (N_{B^0} + N_{\bar{B}^0})$, where N_{B^0} ($N_{\bar{B}^0}$) is the number of B_{tag} mesons identified as a B^0 (\bar{B}^0) for a given measured value of Δt . The signal-to-background ratio is enhanced with a cut on the likelihood ratio as in Fig. 1.

given in Table II. We successfully verified our value of f_{even} with a different method [17] that uses the event rates in $B^+ \rightarrow K^+K_s^0K_s^0$ and the isospin-related channel $B^0 \rightarrow K^+K^-K_s^0$.

To summarize, in a sample of 227 million $B\bar{B}$ meson pairs, we measure the CP content and CP parameters in B^0 -meson decays into ϕK^0 , and into $K^+K^-K_s^0$ with the ϕ mass region excluded. We determine the fraction of CP -even and CP -odd contributions with an angular analysis. In $B^0 \rightarrow \phi K^0$, our values for $\sin 2\beta_{\text{eff}}$ and $C_{\phi K}$ are in good agreement with our previously published val-




FIG. 3: Distributions of S - and P -wave intensities and CP even fraction as a function of K^+K^- invariant mass. Notice that the first bin integrates a wider mass range than the ϕ resonance occupies. Insets show S - and P -wave intensities in the ϕ mass region.

ues [8], and the small S -wave contamination is treated as a systematic uncertainty. In $B^0 \rightarrow K^+K^-K_s^0$, the K^+K^- system is observed to be dominated by S -wave; this, along with the measured value of $\sin 2\beta_{\text{eff}}$, is consistent with previous measurements based on isospin symmetry [9, 10]. Both of our $\sin 2\beta_{\text{eff}}$ values are consistent to within one standard deviation with the value of $\sin 2\beta$ measured in $B^0 \rightarrow c\bar{c}s$ decays [6].

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues, and for the substantial dedicated effort from the computing organizations that support *BABAR*. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF and DFG (Germany), INFN (Italy), FOM (The Netherlands), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from CONACyT (Mex-

ico), A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

* Also with Università della Basilicata, Potenza, Italy

† Deceased

- [1] Throughout this paper, charge conjugate reactions are included implicitly.
- [2] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
- [3] D. London and R. D. Peccei, Phys. Lett. B **223**, 257 (1989); N. G. Deshpande and J. Trampetic, Phys. Rev. D **41**, 895 (1990); R. Fleischer, Z. Phys. C **62**, 81 (1994); N. G. Deshpande and X. G. He, Phys. Lett. B **336**, 471 (1994); Y. Grossman, Z. Ligeti, Y. Nir and H. Quinn, Phys. Rev. D **68**, 015004 (2003); M. Gronau and J. L. Rosner, Phys. Lett. B **564**, 90 (2003).
- [4] A.B. Carter and A.I. Sanda, Phys. Rev. D **23**, 1567 (1981); I.I. Bigi and A.I. Sanda, Nucl. Phys. B **193**, 85 (1981); R. Fleischer and T. Mannel, Phys. Lett. B **511**, 240 (2001); Y. Grossman, G. Isidori and M.P. Worah, Phys. Rev. D **58**, 057504 (1998); Y. Grossman, Z. Ligeti, Y. Nir and H. Quinn, Phys. Rev. D **68**, 015004 (2003); Y. Grossman and M.P. Worah, Phys. Lett. B **395**, 241 (1997); R. Fleischer, Int. J. Mod. Phys. A **12**, 2459 (1997); D. London and A. Soni, Phys. Lett. B **407**, 61 (1997).
- [5] Throughout this paper, ϕ refers to the $\phi(1020)$.
- [6] *BABAR* Collaboration, B. Aubert *et al.*, submitted to Phys. Rev. Lett, hep-ex/0408127.
- [7] Belle Collaboration, K. Abe *et al.*, submitted to Phys. Rev. D, hep-ex/0408111.
- [8] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **93**, 071801 (2004).
- [9] Belle Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **91**, 261602 (2003).
- [10] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **93**, 181805 (2004).
- [11] G. Costa *et al.*, Nucl. Phys. B **175**, 402 (1980); S.U. Chung, Phys. Rev. D **56**, 7299 (1997).
- [12] *BABAR* Collaboration, B. Aubert *et al.*, Nucl. Instr. Methods Phys. Res., Sect. A **479**, 1 (2002).
- [13] Particle Data Group, S. Eidelman *et al.*, Phys. Lett. B **592**, 1 (2004).
- [14] M. Pivk and F. R. Le Diberder, physics/0402083 (2004).
- [15] O. Long, M. Baak, R. N. Cahn, and D. Kirkby, Phys. Rev. D **68**, 034010 (2003).
- [16] *BABAR* Collaboration, B. Aubert *et al.*, Phys. Rev. D **69**, 011102 (2004).
- [17] Belle Collaboration, A. Garmash *et al.*, Phys. Rev. D **69**, 012001 (2004).